

AD-A137 181

SHEATH IONIZATION MODEL OF BEAM EMISSIONS FROM LARGE
SPACECRAFTS(U) AIR FORCE GEOPHYSICS LAB HANSCOM AFB MA
S T LAI ET AL. 30 DEC 83 AFGL-TR-83-0331

1/1

UNCLASSIFIED

F/G 22/1

NL




END

DATE

FILMED

3 SEP 84

DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD A137181

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AFGL-TR-83-0331	2. GOVT ACCESSION NO. AD-A137181	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Sheath Ionization Model of Beam Emissions From Large Spacecrafts		5. TYPE OF REPORT & PERIOD COVERED REPRINT
7. AUTHOR(s) S.T. Lai K. H. Bhavnani* H.A. Cohen M. Tautz*		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Air Force Geophysics Laboratory (PHK) Hanscom AFB Massachusetts 01731		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS 62101F 76611401
11. CONTROLLING OFFICE NAME AND ADDRESS Air Force Geophysics Laboratory (PHK) Hanscom AFB Massachusetts 01731		12. REPORT DATE 30 December 1983
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES 13
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of abstract entered in Block 20, if different from Report) JAN 24 1984		
18. SUPPLEMENTARY NOTES *Radex, Inc., Carlisle, MA Reprinted from Spacecraft Environmental Interactions Technology Conference, USAF Academy, Colorado Springs, CO, 4-6 October 1983		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Sheath Plasma Electron beam Space charge Ionization Ionosphere Spacecraft charging Large spacecraft		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) An analytical model of the charging of a spacecraft emitting electron and ion beams has been applied to the case of large spacecraft. In this model, ionization occurs in the sheath due to the return current. Charge neutralization of spherical space charge flow is examined by solving analytical equations numerically. Parametric studies of potential of large spacecraft are performed. As in the case of small spacecraft, the ions created in the sheath by the returning current play a large role in determining spacecraft potential.		

DTIC FILE COPY

DD FORM 1 JAN 73 1473

84 01 24 004

AFGL-TR-83-0331

SPACECRAFT ENVIRONMENTAL INTERACTIONS TECHNOLOGY CONFERENCE

U.S. AIR FORCE ACADEMY, COLORADO SPRINGS, COLORADO

4-6 October 1983

SHEATH IONIZATION MODEL OF BEAM EMISSIONS FROM LARGE SPACECRAFTS

S. T. Lai and H. A. Cohen
Air Force Geophysics Laboratory, Hanscom AFB., MA.

K. H. Bhavnani and M. Tautz
Radex, Inc., Carlisle, MA.

SUMMARY

An analytical model of the charging of a spacecraft emitting electron and ion beams has been applied to the case of large spacecraft. In this model, ionization occurs in the sheath due to the return current. Charge neutralization of spherical space charge flow is examined by solving analytical equations numerically. Parametric studies of potential of large spacecraft are performed. As in the case of small spacecraft, the ions created in the sheath by the returning current play a large role in determining spacecraft potential.

INTRODUCTION

The potential difference created between spacecraft ground and the ambient plasma during the ejection of a beam of electrons from a sounding rocket payload in the ionosphere (Ref.1) has been found to be much less than had originally been theoretically predicted (Ref.2). To determine the reasons for this limited potential difference, large vacuum chamber tests were conducted in which electron and ion currents were ejected from a payload into a simulated ionosphere.

As a plausible explanation to the observed current voltage behavior, sheath ionization models (Refs. 3,4) for small spacecrafts have been studied. When an electron beam is emitted from a spacecraft, ambient electrons are attracted by the charged spacecraft (Ref.5). They collide with the neutral atmospheric molecules in their paths, and may be energetic enough to ionize the neutrals to form new electrons and ions (Ref. 6). These newly created charges alter the space charge current arriving at the spacecraft shifting the potential to a lower value. The beam electrons are assumed to be energetic enough to leave the spacecraft completely and to play a negligible role in the ionization. This mechanism is capable of explaining the non-monotonic current-voltage behavior observed.

In this paper, we apply the sheath ionization model to the case of large spacecrafts in the ionosphere. In particular, it is important to find out whether the non-monotonic current-voltage behavior during electron beam emissions would still be present for large spacecrafts. Details of the method are given and followed by a discussions of results.

SYMBOLS

e	electron charge
E	electric field
I_b	beam current
m_e	mass of electron
m_i	mass of ion
n_e	density of ambient electrons
n^+	density of ionization ions
n^-	density of ionization electrons
P	probability of ionization
R	radius of spacecraft
r	radial position measured from the center of spacecraft
r'	radial position used as integration variable
r_0	radius of sheath measured from the center of spacecraft
v_e	velocity of an electron in the sheath
v_{th}	thermal velocity of an ambient electron
v_s	sweep velocity
λ	mean free path of electron neutral collision
ϕ	electric potential
ρ	space charge density
ϵ_0	permittivity of space



Form For	
FORM	<input checked="" type="checkbox"/>
718	<input type="checkbox"/>
Uncead	<input type="checkbox"/>
Location	
Distribution/	
Availability Codes	
Avail and/or	
1st	Special
A-1	

MATHEMATICAL FORMULATION

The method of approach used is to study an analytical 'plasma probe' model (Refs. 7,8), with space charge flow of electrons accelerating through the sheath surrounding a spherical 'probe', which represents a spacecraft in an ionizable plasma environment. Magnetic field effect is ignored in this model.

The beam is assumed to be energetic enough to leave the spacecraft completely, and is not stopped by its own space charge at all. As the beam electrons leave, the spacecraft becomes charged oppositely. A polarization region (sheath) is formed in the vicinity of the spacecraft. In our model, ions are assumed depleted due to charge repulsion inside the sheath (Figure 1).

The depletion radius r_0 will be defined by the balance of the outgoing beam current with the incoming ambient current. For a beam current I_b , the depletion radius r_0 is determined by

$$I_b = 4 \pi r_0^2 n_e e v_{th} \quad (1)$$

where v_{th} is the thermal velocity, and n_e is the number density of ambient electrons. Some typical values of sheath radius as calculated by means of eq(1) are shown in figure 2.

The potential ϕ at any point inside the sheath is governed by Poisson's equation :

$$\nabla^2 \phi = - \frac{\rho}{\epsilon_0} \quad (2)$$

where ρ is the space charge density, and ϵ_0 is the permittivity of empty space.

Spherical Symmetric System

To simplify the geometry, we assume spherical symmetry in the spacecraft and sheath system. Equation (2) becomes simply a radial equation :

$$\frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial \phi(r)}{\partial r} \right) = - \frac{\rho(r)}{\epsilon_0} \quad (3)$$

where the gradient of the potential ϕ gives the electric field E :

$$\frac{\partial \phi(r)}{\partial r} = - E(r) \quad (4)$$

Taking into account the electron and ion pairs created as a result of ionization, the charge density ρ at any point r in the sheath is given by the sum of charge densities (figure 3):

$$\rho(r) = e [n^+(r) - n^-(r) - n_e(r)] \quad (5)$$

where n_e is the return current (primary) electron density, while n^+ and n^- are the ionization ion and electron densities respectively, due to return current electron collisions with neutrals.

The ionization electron density $n^-(r)$ is due to all ionizations that occur outwards of r , and the density $n^+(r)$ of ions at r is due to all ionizations that occur inwards of r . Thus,

$$n^-(r) = \frac{1}{r^2} \int_r^{r_0} \frac{\left| \frac{dn}{dt} \right|_{r'} r'^2 dr'}{[2e | \phi(r) - \phi(r') | / m_e]^{1/2}} \quad (6)$$

and

$$n^+(r) = \frac{1}{r^2} \int_R^r \frac{\left[\frac{dn}{dt} \right]_{r'} r'^2 dr'}{[2e |\phi(r) - \phi(r')| / m_i]^{1/2}} \quad (7)$$

where

$$\left[\frac{dn}{dt} \right]_{r'} = \lambda^{-1} P[v_e(r')] n_e(r') v_e(r') \quad (8)$$

Numerical Method

To solve the system of equations (3) to (8), one divides the space of the sheath into N concentric shells, and sets up N equations for the N unknowns ϕ_i (see figure 4). In view of the complexity of the ionization terms in equations (6) and (7), it is impossible to solve these equations exactly. Instead, one seeks the approximate solutions that minimize a function F , the mean square of f_i , constructed from the radial Poisson equation (eq.3) for the i -th cell, where $i=1, \dots, N$.

$$f_i(E_1, \dots, E_N) = (r^2 E)_{i+1} - (r^2 E)_i - \frac{1}{\epsilon_0} (r^2 \rho(E_1, \dots, E_N))_i \Delta r \quad (9)$$

where the electric field E (eq.4) is constructed in a finite difference scheme :

$$\phi_i - \phi_{i+1} = \Delta r (E_i + 2E_{i+1} + E_{i+2})/4 \quad (10)$$

The numerical method used to solve equations (8) to (10) is the standard Newton-Raphson method of iteration :

$$E_1(j+1) = E_1(j) - \frac{f_1(E_1(j), \dots, E_N(j))}{\partial f_1(E_1(j), \dots, E_N(j)) / \partial E_1} \quad (11)$$

A set of trial solutions is used to start in the Newton-Raphson iteration process, and a convergent set of solutions is sought, for each set of input parameters such as beam current, ambient electron density, ambient electron temperature, mean free path, and spacecraft radius.

RESULTS AND DISCUSSION

Figure 5 shows the computed results of spacecraft potential as a function of electron beam current for various electron density, electron temperature, and mean free path. The non-monotonic behavior of potential current curves shows up. At low currents, the potential increases with beam current. When the current increases further, ionization occurs inside the sheath. The potential then turns around as the current of the electron beam increases.

The ion and electron charges created by ionization alter the behavior of the space charge flow, originally governed by the single charged Poisson equation. The potential turns to a lower value, and stays approximately constant as current further increases.

At this stage, the potential profile as a function of radial distance show locally flat gradient. This is due to ions created inside the sheath not being able to move out quickly due to their heavy masses. If a local ion charge build-up forms a potential hump, ion motion would be twoways, and, the theory would then break down.

To overcome this difficulty, a sweep velocity v_s is added to the ions. Eq.(7) becomes

$$n^+(r) = \frac{1}{r^2} \int_R^r \frac{\left[\frac{dn}{dt} \right]_{r'} r'^2 dr'}{[2e|\phi(r) - \phi(r')|/m_i + v_s^2]^{1/2}} \quad (12)$$

It is argued that the motion of a spacecraft relative to its plasma environment can provide such a sweep velocity v_s (eq.(12)). The value of v_s is of the order of spacecraft velocity and is an arbitrary input to the computation. However, at a higher current, a potential hump again shows up, and the computation fails to converge. The technique breaks down. It is conjectured that twoway space charge flows should be accommodated when a potential hump appears.

For increasing spacecraft radii, the non-monotonic current-voltage behavior still persists (see figure 6). However, increased spacecraft radius lowers the maximum spacecraft potential induced by beam emission. Also, the amplitude of the difference between the maximum potential and the minimum (beyond the turn-around) diminishes. Figure 7 shows a plot of the envelope of maximum and minimum potentials for various spacecraft radii.

For a given beam current I_b (see eq.(1)), the sheath surface area remains constant and is unaffected by the increase in spacecraft radius. The sheath thickness (defined as the sheath radius minus the spacecraft radius), however, diminishes. As a result, a lower spacecraft potential is sufficient to attract ambient electrons, through the sheath, for the compensation of electron beam current leaving the spacecraft.

Beyond the turn-around point in a current-potential curve, the minimum potential is limited by the minimum energy required to ionize a neutral molecule in the atmosphere. Since such a minimum energy is generally of the order of 20 eV. (Ref.6), the minimum potential in a current-potential curve is expected to approach about 20 eV. asymptotically, depending on the model of ionization used. For the same reason, if the maximum potential induced by beam emissions is below about 20 eV., no non-monotonic behavior is expected.

Figure 7 shows the calculated envelopes of the maximum and minimum (beyond turn-around) potentials for various spacecraft radii, in a given ambient environment. The amount of ionization becomes very small as the sheath potential approaches the minimum ionization potential. The amplitude of the potential drop beyond the turn-around also approaches the value of minimum ionization energy.

There is another critical beam current, which manifests itself for large spacecrafts, but not for small ones. This current is determined by equating the sheath radius to the spacecraft radius. If the sheath radius is too small, the spacecraft would receive enough ambient electrons to

compensate beam emissions without being charged up. The potential of the spacecraft is that of natural charging, in this case. Beyond this critical current, the beam emission is able to swing the spacecraft to an opposite potential and hence control the charging of the spacecraft. This phenomenon shows up in the calculations (see figure 6).

In the model studied, as the radius of a spacecraft increases, three regimes of physical behavior can be identified. Figure 8 shows these regimes clearly. The potential versus spacecraft radius curve is relatively flat in the small radius regime. This is the regime in which saturated ionization occurs, i.e. this is the regime beyond the minimum potential in a current-voltage curve. The second regime is characterized by the presence of the potential maximum, which is the main feature of non-monotonic behavior. The third regime occurs when the spacecraft is so large that its radius exceeds the sheath radius (measured from the spacecraft center) for a given current. The beam loses its control of the spacecraft potential, and natural charging dominates.

REFERENCES

1. O'Neil R.R., Bien F., Burt D., Sandock J.A., and Stair A.T. Jr.: Summarized Results of the Artificial Auroral Experiment PRECEDE. J. Geophys. Res. Vol.83, No.A7, p.3273, 1978.
2. Parker L.W. and Murphy B.L.: Potential Buildup on an Electron-Emitting Ionospheric Satellite. J. Geophys. Res. Vol.72, p.1631, 1967.
3. Leadon R.E., Woods A.J., Wenaas E.P., Klein H.H.: Analytical Investigation of Emitting Probes in an Ionized Plasma. AFGL-TR-81-0138, 1981; ADA104166.
4. Cohen H.A., Lai S.T., McNeil W.J., Wenaas E.P., Leadon R.E.: Spacecraft Charging with Beam Emissions in an Ionizable Environment. EOS Vol.64, No.18, p.301, 1983.
5. Beard D.B. and Johnson F.S.: Ionospheric Limitations on Attainable Satellite Potential. J. Geophys. Res. Vol.66, pp.4113-4122. 1961.
6. Rapp D., Englander-Golden P.: Total Cross-Sections for Ionization and Attachment by Electron Impact in Gases. J. Chem. Phys. Vol.43, p.1464, 1965.
7. Chen F.F.: Electric Probes, in Plasma Diagnostic Techniques, ed. by Huddleston R.H. and Leonard S.L., Academic Press. 1965.
8. Lam S.H.: Unified Theory of the Langmuir Probe in a Collisionless Plasma, Phys. Fluids. Vol.8, No.1, p.73, 1965.




Figure 1. Sheath Formation during Beam Emission.




Figure 2. Parametric Dependence of Sheath Size




Figure 3. Ionization Pair-Creation in Sheath




Figure 4. Decomposition of Sheath into Shells




Figure 5. Relation Between Potential Profile and I-V Behavior.




Figure 6. Persistence of Non-monotonic I-V Behavior. The Parametric Conditions are as in Figure 7.




Figure 7. Envelope of the Potential Extrema in Figure 6.




Figure 8. Non-monotonic Behavior of Spacecraft Potential as a Function of Spacecraft Radius, for a Given Electron Beam Current.

DA
FILM